



**Lunch and Learn Series
IC and Ultrasound
Reference & Resource List**

Toolkit

Ultrasound Infection Prevention Toolkit <https://www.ultrasoundinfectionprevention.org/>

CDC

Guideline for Disinfection and Sterilization in Healthcare Facilities (2008. Updated in 2017) Retrieved from <https://www.cdc.gov/infectioncontrol/guidelines/disinfection/index.html>

FDA

FDA (2019). Marketing Clearance of Diagnostic Ultrasound Systems <https://www.fda.gov/media/71100/download>

FDA. Ultrasound Imaging. Available from <https://www.fda.gov/radiation-emitting-products/medical-imaging/ultrasound-imaging>

Articles

Society off Diagnostic Medical Sonography. (2020). Guidelines for Infection Prevention and Control in Sonography: Reprocessing the Ultrasound Transducer <https://www.sdms.org/docs/default-source/Resources/8756479320933256.pdf>

Scott, D., Fletcher, E., Kane, H., Malcolm, W., Kavanagh, K., Banks, A. L., & Rankin, A. (2018). Risk of infection following semi-invasive ultrasound procedures in Scotland, 2010 to 2016: A retrospective cohort study using linked national datasets. *Ultrasound*, 26(3), 168-177. doi: 10.1177/1742271X18774594

Koibuchi, H., Kotani, K., & Taniguchi, N. (2013). Ultrasound probes as a possible vector of bacterial transmission. *Medical ultrasonography*, 15(1), 41-44. doi: 10.11152/mu.2013.2066.151.hk1upp2

Casalegno, J. S., Carval, K. L. B., Eibach, D., Valdeyron, M. L., Lamblin, G., Jacquemoud, H., ... & Mekki, Y. (2012). High risk HPV contamination of endocavity vaginal ultrasound probes: an underestimated route of nosocomial infection?. *PloS one*, 7(10), e48137. doi: 10.1371/journal.pone.0048137

Carrico, R. M., Furmanek, S., & English, C. (2018). Ultrasound probe use and reprocessing: Results from a national survey among US infection preventionists. *American journal of infection control*, 46(8), 913-920. doi: 10.1016/j.ajic.2018.03.025

Carrico, R.. (2019) Ultrasound infection prevention: Risks, challenges and solutions [PowerPoint]. Retrieved from https://apicchicago.starchapter.com/images/downloads/Spring_2019_3M_Conference/ultrasound_infection_prevention_risks_challenges_and_solutions.pdf

Westerway, S. C., Basseal, J. M., Brockway, A., Hyett, J. A., & Carter, D. A. (2017). Potential infection control risks associated with ultrasound equipment—a bacterial perspective. *Ultrasound in medicine & biology*, 43(2), 421-426. doi: 10.1016/j.ultrasmedbio.2016.09.004

Gillespie, J. L., Arnold, K. E., Noble-Wang, J., Jensen, B., Arduino, M., Hageman, J., & Srinivasan, A. (2007). Outbreak of *Pseudomonas aeruginosa* infections after transrectal ultrasound-guided prostate biopsy. *Urology*, 69(5), 912-914. doi: 10.1016/j.urology.2007.01.047

Nyhsen, C. M., Humphreys, H., Koerner, R. J., Grenier, N., Brady, A., Sidhu, P., ... & Claudon, M. (2017). Infection prevention and control in ultrasound—best practice recommendations from the European Society of Radiology Ultrasound Working Group. *Insights into imaging*, 8(6), 523-535. doi: 10.1007/s13244-017-0580-3

Reese SM, et al. (2019). Implementation of cleaning and evaluation process for mobile patient equipment using adenosine triphosphate. *Infection Control & Hospital Epidemiology*, 40: 798–800. doi.org/10.1017/ice.2019.115

Ai, A., Anderson, L., Safdar, N., Shirley, D. (2020). Barriers and facilitators to standardization of ultrasound use and probe disinfection in the ambulatory setting. *Infection Control & Hospital Epidemiology*, 41(4) pp. 469 – 471. doi:10.1017/ice.2020.13

Gaillot, O., Maruéjols, C., Abachin, É., Lecuru, F., Arlet, G., Simonet, M., & Berche, P. (1998). Nosocomial outbreak of *Klebsiella pneumoniae* producing SHV-5 extended-spectrum β -lactamase, originating from a contaminated ultrasonography coupling gel. *Journal of clinical microbiology*, 36(5), 1357-1360. doi:10.1128/JCM.36.5.1357-1360.1998

Shaban, R. Z., Maloney, S., Gerrard, J., Collignon, P., Macbeth, D., Cruickshank, M., ... & Derrington, P. (2017). Outbreak of health care-associated *Burkholderia cenocepacia* bacteremia and infection attributed to contaminated sterile gel used for central line insertion under ultrasound guidance and other procedures. *American journal of infection control*, 45(9), 954-958. doi: 10.1016/j.ajic.2017.06.025

Infection prevention and control in ultrasound—best practice recommendations from the European Society of Radiology Ultrasound Working Group. *Insights into imaging*, 8(6), 523-535. doi:10.1007/s13244-017-0580-3

Ryndock, E., Robison, R., Meyers, C. (2016) Susceptibility of HPV16 and 18 to high level disinfectants indicated for semi-critical ultrasound probes. *J Medical Virology*, 88(6):1076–1080. doi: 10.1002/jmv.24421

Meyers J., Ryndock, E., Conway, M., J., Meyers, C., Robison, R. (2014) Susceptibility of high-risk human papillomavirus type 16 to clinical disinfectants. *J Antimicrob Chemotherapy*, 69(6):1546–1550. doi: 10.1093/jac/dku006

Oleszkowicz, S. C., Chittick, P., Russo, V., Keller, P., Sims, M., & Band, J. (2012). Infections associated with use of ultrasound transmission gel: proposed guidelines to minimize risk. *Infect Control Hosp Epidemiol*, 33(12), 1235-1237. doi: 10.1086/668430